


## WHAT IS DE QUERVAIN'S STENOSING TENOSYNOVITIS?


Once called Washer Woman's Sprain, this painful inflammation of the thumb side of the wrist now bears the name of the Swiss surgeon, Fritz de Quervain, who wrote about this condition in 1895.

## ANATOMY

Passing over the back (or dorsal surface) of the wrist are the tendons for muscles that extend or straighten the fingers and thumb, and lift the hand at the wrist. These tendons run through six lubricated tunnels (compartments) under a thick fibrous layer called the extensor retinaculum.



**The first dorsal compartment** lies over the bony bump at the base of the thumb. Through it pass the tendons for the abductor pollicis longus (APL) and the extensor pollicis brevis (EPB) muscles. Both of these muscles help spread and extend the thumb away from the rest of the hand.



*Anatomical position of the first dorsal compartment*

## WHAT IS IT?

De Quervain's Stenosing Tenosynovitis is a painful inflammation of the tendons in the first dorsal compartment of the wrist. The lubricating synovial sheath lining this compartment thickens and swells, giving the enclosed tendons less room to move, and produces extra synovial fluid. A painful cyst may also form.

## WHAT CAUSES IT?

This inflammation may be caused by anything that changes the shape of the compartment or causes swelling or thickening of the tendons.


## WHO GETS IT?

De Quervain's Stenosing Tenosynovitis occurs most often in individuals between the ages of 30 and 50. Women are afflicted with this condition 8 to 10 times more often than men. People who engage in activities requiring sideways motion of the wrist while gripping the thumb, as in hammering, skiing, and some assembly line jobs, may be predisposed to developing this disorder.

## WHAT ARE THE SYMPTOMS?

Pain over the thumb side of the wrist is the primary symptom. This condition may occur "overnight" or very gradually, and it may radiate pain into the thumb and up the forearm. It is worse with the use of the hand and thumb, especially with any forceful grasping, pinching, or twisting. Swelling over the thumb side of the wrist may be present, as well as some "snapping" when the thumb is moved. Because of the pain and swelling, there may be some decreased thumb motion.

Besides pain and swelling over the first dorsal compartment, having a positive Finkelstein's Test is a good indication the patient has this problem. In this test, the patient makes a fist with his or her thumb placed under the fingers. Then the patient bends the wrist away from the thumb and towards the little finger side of the hand. This test is mildly painful to many of us, but to someone with de Quervain's Stenosing Tenosynovitis, it is very painful.




## CONSERVATIVE MANAGEMENT

Your doctor may first try to reverse the course of the disease with a 3 to 6 week trial of anti-inflammatory medication, while the wrist and thumb are rested by wearing a wrist and thumb spica splint. Your physician may also inject the area with a steroid to help decrease the inflammation.

## IF SURGERY IS NECESSARY

If the symptoms of de Quervain's Stenosing Tenosynovitis disease are long-standing or unresponsive to conservative management, surgery is indicated. This type of surgery is usually performed on an outpatient basis.

During surgery, an incision is made over the first dorsal compartment. The abnormally thickened dorsal carpal ligament is cut to release the APL and EPB tendons. The wound is then closed, and a compressive dressing is applied.



*Wrist and thumb splint/bulky dressing*

## POST-OP

A prescription for pain medication will be provided. The bulky dressing is removed within 1 to 2 weeks after surgery, and an exercise program for the thumb and wrist is started. It usually takes several weeks for a full recovery. DeQuervain's Stenosing Tenosynovitis rarely recurs after proper surgical intervention. Complications may occur if not treated.

# DE QUERVAIN'S STENOSING TENOSYNOVITIS



The Louisiana Hand  
and Elbow Center

T (985) 625-2200  
F (985) 625-2206

604 North Acadia Road, Suite 508  
Thibodaux, LA 70301



The Louisiana Hand  
and Elbow Center